

A Strong Lyman- α Emitter at $z = 6.33$ in the Subaru Deep Field Selected as an i' Dropout¹

T. Nagao ^{2,3}, Y. Taniguchi ³, N. Kashikawa ⁴, K. Kodaira ⁵, N. Kaifu ⁴, H. Ando ⁴, H. Karoji ⁶, M. Ajiki ³, M. Akiyama ⁶, K. Aoki ⁶, M. Doi ⁷, S. S. Fujita ³, H. Furusawa ⁶, T. Hayashino ⁸, F. Iwamuro ⁹, M. Iye ⁴, N. Kobayashi ⁷, T. Kodama ⁴, Y. Komiyama ⁶, Y. Matsuda ^{4,8}, S. Miyazaki ⁶, Y. Mizumoto ⁴, T. Morokuma ⁷, K. Motohara ⁷, T. Murayama ³, K. Nariai ¹⁰, K. Ohta ⁹, S. Okamura ^{11,12}, M. Ouchi ⁴, T. Sasaki ⁶, Y. Sato ⁴, K. Sekiguchi ⁶, K. Shimasaku ¹¹, Y. Shioya ³, H. Tamura ⁸, I. Tanaka ³, M. Umemura ¹³, T. Yamada ⁴, N. Yasuda ¹⁴, & M. Yoshida ¹¹

ABSTRACT

We report on the discovery of a star-forming galaxy at $z = 6.33$ in the Subaru Deep Field. This object is selected as a candidate of an i' -dropout, high-redshift

¹Based on data collected at the Subaru Telescope, which is operated by the National Astronomical Observatory of Japan.

²INAF – Osservatorio Astrofisico di Arcetri, Largo Enrico Fermi 5, 50125 Firenze, Italy; tohru@arcetri.astro.it

³Astronomical Institute, Graduate School of Science, Tohoku University, Aramaki, Aoba, Sendai 980-8578, Japan

⁴National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan

⁵The Graduate University for Advanced Studies, Shonan Village, Hayama, Kanagawa 240-0193, Japan

⁶Subaru Telescope, National Astronomical Observatory of Japan, 650 N. A'ohoku Place, Hilo, HI 96720

⁷Institute of Astronomy, Graduate School of Science, University of Tokyo, 2-21-1 Osawa, Mitaka, Tokyo 181-0015, Japan

⁸Research Center for Neutrino Science, Graduate School of Science, Tohoku University, Aramaki, Aoba, Sendai 980-8578, Japan

⁹Department of Astronomy, Graduate School of Science, Kyoto University, Kitashirakawa, Sakyo, Kyoto 606-8502, Japan

¹⁰Department of Physics, Meisei University, 2-1-1 Hodokubo, Hino, Tokyo 191-8506, Japan

¹¹Department of Astronomy, Graduate School of Science, University of Tokyo, Tokyo 113-0033, Japan

¹²Research Center for the Early Universe, Graduate School of Science, University of Tokyo, Tokyo 113-0033, Japan

¹³Center for Computational Physics, University of Tsukuba, 1-1-1 Tennodai, Tsukuba 305-8571, Japan

¹⁴Institute for Cosmic Ray Research, University of Tokyo, Kashiwa 277-8582, Japan

galaxy around $z = 6$ because of its red $i' - z'$ color in our deep optical imaging survey in the Subaru Deep Field. Our follow up optical spectroscopy reveals that this object is a strong Ly α emitter with only very faint ultraviolet continuum. The rest-frame equivalent width of the detected Ly α emission is as much as 130 Å. Thus the light detected in our z' image is largely attributed to the Ly α emission, i.e., $\sim 40\%$ of the z' -band flux is the strong Ly α emission, giving a very red $i' - z'$ color. This is consistent with the photometric property of this object because the narrow-band data obtained with the NB921 filter shows a significant depression, $z' - \text{NB921} = -0.54$ mag. By using the photometric data, we show that some other objects among the 48 i' -dropout high-redshift galaxy candidates found in the Subaru Deep Field also show a significant NB921 depression. We briefly discuss the nature of these NB921-depressed objects.

Subject headings: early universe — galaxies: evolution — galaxies: formation — galaxies: individual (SDF J132440.6+273607) — galaxies: starburst

1. INTRODUCTION

The star formation in the early universe provides important clues on the understanding of both galaxy formation and the ionization state of intergalactic matter (e.g., Loeb & Barkana 2001). Currently, two alternative methods are frequently used to find very high- z objects; one is to search for strong emission-line objects by using narrow-passband filters (e.g., Hu et al. 2002; Ouchi et al. 2003; Ajiki et al. 2003; Kodaira et al. 2003; see for a review, Taniguchi et al. 2003), which selectively picks up high- z objects with a strong emission line such as Ly α (Ly α emitters; hereafter LAEs). The other method is to search for redshifted Lyman-break objects based on broad-band color selections (e.g., Steidel et al. 1996a, 1996b; 1999; Iwata et al. 2003; Stanway, Bunker, & McMahon 2003; Ouchi et al. 2004; Giavalisco et al. 2004). Since this method relies on the stellar Lyman-break spectral feature, this method tends to select high- z galaxies with strong UV continuum emission, i.e., Lyman-break galaxies (LBGs). Since some of LBGs also show a strong Ly α emission, LAEs may be regarded as a subclass of LBGs (see, e.g., Shapley et al. 2003).

And very recently, based on imaging surveys with broad pass-band filters, objects with a very red $i' - z'$ color (i.e., i' dropout objects) are intensively explored as candidates of galaxies at $z \gtrsim 6$, where the cosmic reionization ended (e.g., Stanway et al. 2003; Bouwens et al. 2003, 2004; Dickinson et al. 2004; Bunker et al. 2004). In this Letter, we report on the discovery of a luminous star-forming galaxy at $z = 6.33$ in the Subaru Deep Field (SDF). This object is originally selected as a candidate of i' -dropout high-redshift galaxy because

of its red $i' - z'$ color by the imaging survey for SDF. This object is interesting because our follow-up spectroscopy shows that it presents only very faint UV continuum emission. We adopt a cosmology with $(\Omega_{\text{tot}}, \Omega_{\text{M}}, \Omega_{\Lambda}) = (1.0, 0.3, 0.7)$ and $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ throughout this Letter. We use the AB photometric system for optical magnitudes.

2. OBSERVATIONS

2.1. Optical Deep Imaging

We have carried out a very deep optical imaging survey in the SDF centered at $\alpha(\text{J2000}) = 13^{\text{h}} 24^{\text{m}} 38^{\text{s}}.9$ and $\delta(\text{J2000}) = +27^\circ 29' 25''.9$, by using Suprime-Cam (Miyazaki et al. 2002) which consists of 5×2 CCDs of $2\text{k} \times 4\text{k}$ pixels with a pixel scale of $0''.202 \text{ pixel}^{-1}$, on the 8.2m Subaru Telescope (Kaifu et al. 2000; Iye et al. 2004). The observations were made with five broad pass-band filters, B , V , R_c , i' , and z' , and two narrow pass-band filters, $NB816$ and $NB921$. The central wavelengths and the half widths of the transmittance of the two narrow pass-band filters are $(\lambda_c, \Delta\lambda_{\text{FWHM}}) = (8150\text{\AA}, 120\text{\AA})$ and $(9196\text{\AA}, 132\text{\AA})$; see Ajiki et al. (2003) and Kodaira et al. (2003) for more details. The data were collected in several observing runs during a period between 2001 and 2003. A summary of the imaging observations is given in Table 1.

The individual CCD data were reduced and combined using IRAF and the mosaic-CCD data reduction software developed by Yagi et al. (2002). The combined images for individual bands were aligned and smoothed with Gaussian kernels to match their seeing sizes. The PSF FWHM of the final images is $0''.98$. Exposure times and limiting magnitudes are listed in Table 1. Photometric calibrations are made using usual standard stars. Source detection and photometry are performed using SExtractor version 2.1.6 (Bertin & Arnouts 1996). Here the z' image is used for the source detection. The effective area of this imaging survey is 761 arcmin^2 .

We select candidates of i' -dropout galaxies at $z \sim 6$ imposing the following four criteria on the z' -selected sample; $i' - z' > 1.5$, $z' < 26.1$ (above 5σ), $B > 28.5$ (below 3σ), and $R_c > 27.8$ (below 3σ), where the magnitudes are measured by adopting the aperture size of a $2''.0$ diameter. We finally obtain a photometric sample of 48 i' -dropout galaxy candidates at $z \sim 6$.

2.2. Optical Spectroscopy

In order to investigate the nature of i' -dropout high- z galaxy candidates found in our optical imaging survey, we have completed optical spectroscopy of 9 objects in our 48 i' -dropout sample up to now. We used the Subaru Faint Object Camera And Spectrograph (FOCAS; Kashikawa et al. 2002) on 2003 May and June. Among the 9 objects, we concentrate our discussion on an interesting object, SDF J132440.6+273607, in this Letter. The observational properties of the other i' -dropout sample will be described elsewhere. For the spectroscopy of SDF J132440.6+273607, the 300 lines mm^{-1} grating was used with an order cut filter O58. The wavelength coverage was $\sim 5800 \text{ \AA}$ to $10000 \text{ \AA}$ with a pixel resolution of $1.35 \text{ \AA}$. The use of a $0''.8$ slit gave a spectroscopic resolution of $9.0 \text{ \AA}$ at $9000 \text{ \AA}$ that is measured by atmospheric OH airglow lines. This corresponds to the spectral resolution of $R \simeq 1000$. The spatial sampling was $0''.31$ per resolution element as we adopted 3 pixel on-chip binning. We obtained seven 1800s exposure frames for SDF J132440.6+273607. Typical seeing size was $0''.4 - 0''.6$ during the observation. We also obtained the spectra of spectroscopic standard stars Hz 44 and Feige 34 for flux calibration.

3. RESULTS

As mentioned above, SDF J132440.6+273607 was selected by its red $i' - z'$ color, i.e., so-called i' -dropout selection. Indeed this object is not detected above the 3σ level on the B , V , R_c , i' , and $NB816$ images. SDF J132440.6+273607 is clearly detected in the z' and $NB921$ images; the $2''.0$ aperture magnitudes are $z' = 25.66$ mag and $NB921 = 26.20$ mag. Since the 3σ limiting magnitude of i' is $i'_{\text{lim}} = 27.43$ mag, SDF J132440.6+273607 has a color of $i' - z' > 1.77$, satisfying our criterion for the i' -dropout selection, $i' - z' > 1.5$. The optical thumbnail images are shown in Figure 1. The image size of SDF J132440.6+273607 in FWHM is 5.8 pixels or $1''.1$ in the z' -band image and 6.0 pixels or $1''.2$ in the $NB921$ -band image, respectively. Both of these two images may be only marginally larger than the PSF size.

In Figure 2, we show the sky-subtracted optical position-velocity spectrogram and sky-subtracted one-dimensional spectrum of SDF J132440.6+273607, obtained by FOCAS on Subaru. The aperture size to extract the one dimensional spectrum is 5 binned pixels ($1''.6$). We can see a prominent single emission line whose peak is at $8909 \text{ \AA}$, and no significant continuum emission in the whole of the covered wavelength range. The spectral feature seen at $\sim 8885 \text{ \AA}$ is a residual of the sky subtraction. The observed velocity width of the emission line is $\sim 8 \text{ \AA}$ in FWHM, roughly comparable to the instrumental width. However, we can see a clear asymmetric feature below the half maximum of the emission line; i.e.,

a very prominent redward tail which extends up to $\sim 8930\text{\AA}$. This asymmetry and the photometric property strongly suggest that the detected emission line is the $\text{Ly}\alpha$ emission and that SDF J132440.6+273607 is at $z = 6.33$. Note that this object is thought not to be an emission-line object at lower redshift; i.e., an $\text{H}\alpha$ emitter at $z \sim 0.36$, an $[\text{O III}]\lambda 5007$ emitter at $z \sim 0.78$, or an $\text{H}\beta$ emitter at $z \sim 0.83$. This is because there is no emission lines in the covered wavelength range other than the prominent emission at $8909\text{\AA}$. We cannot completely deny the possibility that SDF J132440.6+273607 is a $[\text{O II}]\lambda 3727$ emitter at $z \sim 1.39$, because the $[\text{O II}]\lambda 3727$ doublet cannot be resolved by our observation (the expected wavelength separation of the $[\text{O II}]$ doublet is $6.7\text{\AA}$). However, this does not seem plausible because the observed red tail or shoulder of the detected emission line is hard to be understood if the emission line is $[\text{O II}]\lambda 3727$. Therefore, taking both the photometric and the spectroscopic properties into account, we conclude that SDF J132440.6+273607 is a strong $\text{Ly}\alpha$ emitter without strong UV stellar continuum emission at $z = 6.33$. The observed $\text{Ly}\alpha$ flux is $(4.0 \pm 0.1) \times 10^{-17} \text{ ergs s}^{-1} \text{ cm}^{-2}$, where the slit loss is not taken into account. The $\text{Ly}\alpha$ luminosity is thus calculated to be $1.8 \times 10^{43} \text{ ergs s}^{-1}$.

Based on the $\text{Ly}\alpha$ flux obtained by the spectroscopy, we can derive a $\text{Ly}\alpha$ -corrected z' magnitude, $z'_{\text{cor}} = 26.16 \text{ mag}$, which is very close to the $NB921$ magnitude. Since our i' -dropout search is complete only down to $z' = 25.93 \text{ mag}$, we could not pick up SDF J132440.6+273607 as an i' -dropout object if there were no strong $\text{Ly}\alpha$ emission; i.e., $i' - z'_{\text{cor}} > 1.27$, which does not satisfy our i' -dropout criterion ($i' - z' > 1.5$). Then we focus on the depressed $NB921$ magnitude compared to the z' magnitude; i.e., $z' - NB921 = -0.54 \text{ mag}$. If we assume that the depression of the $NB921$ magnitude is caused only by the contribution of the $\text{Ly}\alpha$ emission in the z' band, the required flux of the $\text{Ly}\alpha$ emission is $(4.2 \pm 2.2) \times 10^{-17} \text{ ergs s}^{-1} \text{ cm}^{-2}$. This is consistent with the spectroscopically measured $\text{Ly}\alpha$ flux, where we assume that all of the $NB921$ flux is attributed to the stellar UV continuum emission. This assumption leads to the UV continuum flux density of $(4.3 \pm 1.0) \times 10^{-20} \text{ ergs s}^{-1} \text{ cm}^{-2} \text{ \AA}^{-1}$, which is consistent with the non-detection of the continuum emission in our optical spectroscopy of SDF J132440.6+273607 (Figure 2).

4. DISCUSSION

As presented in the last section, our spectroscopic observation discloses that SDF J132440.6+273607 is a strong LAE with only faint UV stellar emission, despite the fact that it has been originally selected as an i' -dropout galaxy candidate by its red $i' - z'$ color. This LAE is thought to be an actively star-forming galaxy, though we cannot completely reject the possibility that it is a narrow-line QSO. Although the stellar continuum emission

is not detected in our spectroscopy, we can estimate the equivalent width of the $\text{Ly}\alpha$ emission based on the $NB921$ data. The photometrically determined $\text{Ly}\alpha$ equivalent width of SDF J132440.6+273607 is $EW_{\text{obs}}(\text{Ly}\alpha) = 980\text{\AA}$. We then obtain the equivalent width at the rest frame; $EW_0(\text{Ly}\alpha) = 130\text{\AA}$. The derived equivalent width is very large as a broad-band selected galaxy, since a very small fraction of LBGs at $z \sim 3$ show $EW_0(\text{Ly}\alpha) > 100\text{\AA}$ (e.g., Steidel et al. 1999; Shapley et al. 2003). Rather this extremely large equivalent width seems more similar to the narrow-band selected galaxies (e.g., Malhotra & Rhoads 2002; Ajiki et al. 2003; Hu et al. 2004). Hu et al. (2004) recently found a number of LAEs with $EW_0(\text{Ly}\alpha) > 100\text{\AA}$ at $z \sim 5.7$ by using a narrow-passband filter. Note that LBGs and LAEs do not necessarily have completely distinct properties such as $EW_0(\text{Ly}\alpha)$, because they are defined just by their selection method. However, it is not clear how these two populations overlap or how different statistically, especially at the very high- z universe where the number of galaxies that have been investigated by spectroscopic observations is very small. Spectroscopic observations on a large number of high- z galaxies are also necessary to investigate whether or not the $EW_0(\text{Ly}\alpha)$ frequency distribution depends on redshift, which is also a very interesting issue. Forthcoming intensive spectroscopic observations for high- z galaxies will hopefully solve these problems.

Our finding of SDF J132440.6+273607 naturally leads to the following question: is this object a very peculiar object, or does any photometrically constructed i' -dropout sample generally contain a lot of galaxies with only faint UV stellar continuum? In order to investigate this issue, we focus on the photometrically-selected 48 i' dropout LBG candidates in SDF. In Figure 3, we show the $z' - NB921$ colors of the i' dropout LBG candidates as functions of the z' magnitude and the $i' - z'_{\text{cor}}$ color. The $NB921$ depression of SDF J132440.6+273607 is $z' - NB921 = -0.54$ as mentioned above, whose statistical significance is roughly $\sim 2\sigma$. Interestingly, there are other three i' -dropout galaxy candidates which show significant $NB921$ depression above the 3σ level (objects b, c, and e in the Figure 3).

There are three possibilities for such $NB921$ -depressed, i' -dropout high- z galaxy candidates. The first possibility is the LAE at $6.0 \lesssim z \lesssim 6.5$ which corresponds to $8510\text{\AA} \lesssim \lambda_{\text{Ly}\alpha} \lesssim 9110\text{\AA}$, just similar to the case of SDF J132440.6+273607. In this redshift range, the $\text{Ly}\alpha$ emission falls only in the z' band and not in the $NB921$ one, which results in the $NB921$ depression. The second possibility is the “ $NB921$ -dropout LBG”; i.e., the LBGs at $z \gtrsim 6.6$. In this case, the original categorization as i' dropout LBGs is not wrong, because the objects show the UV stellar continuum with a significant Lyman break. The third possibility is the i' dropout galaxy with a significant absorption feature at the $NB921$ range, although it is not clear whether or not such an object really exists in such a high-redshift universe. In order to investigate these ideas further, we plot the expected colors of high- z galaxies with a strong $\text{Ly}\alpha$ emission on the $z' - NB921$ versus $i' - z'$ two-color diagram (Figure 3).

Here we use the galaxy evolution model of Bruzual & Charlot (2003) by assuming that the star-formation rate of model galaxies is proportional to $\exp(-t/\tau)$, where $\tau = 1$ Gyr and $t = 1$ Gyr are adopted. Here we examine only three cases, i.e., $EW_0(\text{Ly}\alpha) = 65, 130$, and $260 \text{ \AA}$. The $NB921$ depression of $-0.2, -0.6$ and -1.0 mag can be realized for the cases of $EW_0(\text{Ly}\alpha) = 65, 130$, and $260 \text{ \AA}$ for the first possibility. On the other hand, more significant $NB921$ depression appears for the second possibility. As for the second possibility, a very red color of $i' - z' > 3$ is also expected. However, an extremely deep i' image is required to discriminate these possibilities clearly. Since the two $NB921$ -depressed i' dropout objects with $i' - z' \sim 3$ may be $NB921$ -dropout galaxies, a follow-up spectroscopic observation will be very interesting to access galaxies at $z \gtrsim 6.6$.

Here we discuss the star-forming activity in SDF J132440.6+273607. The $\text{Ly}\alpha$ flux of this object measured by our spectroscopy is $4.0 \times 10^{-17} \text{ ergs s}^{-1} \text{ cm}^{-2}$, which corresponds to the $\text{Ly}\alpha$ luminosity of $1.8 \times 10^{43} \text{ ergs s}^{-1}$. We can estimate the star-formation rate (SFR) by using the following relation; $SFR(\text{Ly}\alpha) = 9.1 \times 10^{-43} L(\text{Ly}\alpha) M_\odot \text{ yr}^{-1}$ (Kennicutt 1998; Brocklehurst 1971). We then obtain $SFR(\text{Ly}\alpha) \simeq 16 M_\odot \text{ yr}^{-1}$. This is a lower limit because no correction was made for possible absorption effects on the $\text{Ly}\alpha$ emission. We can also estimate the SFR by the luminosity of the UV stellar continuum by adopting the following relation; $SFR(\text{UV}) = 1.4 \times 10^{-28} L_\nu M_\odot \text{ yr}^{-1}$ (Kennicutt 1998). By using the UV continuum flux density measured from the $NB921$ magnitude, $f_\nu(NB921) = 1.2 \times 10^{-30} \text{ ergs s}^{-1} \text{ cm}^{-2} \text{ Hz}^{-1}$, we obtain $SFR(\text{UV}_{1255}) \simeq 10.4 M_\odot \text{ yr}^{-1}$. One of the reason why $SFR(\text{UV}_{1255})$ is smaller than $SFR(\text{Ly}\alpha)$ may be that $SFR(\text{UV})$ would be underpredicted for objects with a very large $EW_0(\text{Ly}\alpha)$. This is because such a huge $EW_0(\text{Ly}\alpha)$ is hard to be created by a continuous star formation with a normal initial-mass function, which is assumed in the above relation between the SFR and the luminosity of the UV stellar continuum. Schaerer (2000) presented that $SFR(\text{UV})$ is underestimated for star-forming galaxies with an age of $< 10^8 \text{ yr}$. Taking all of above considerations into account, we conclude that the SFR of SDF J132440.6+273607 is $> 16 M_\odot \text{ yr}^{-1}$. Since there may be a large number of objects with large $EW_0(\text{Ly}\alpha)$ as suggested by Figure 3, systematic spectroscopic studies on i' -dropout galaxies are crucial to understand the star-forming activities of galaxies at $z \gtrsim 6$, which is a very important epoch in relation to the cosmic reionization.

We thank the referee, A. Bunker, for his valuable comments. We also thank the Subaru Telescope staff for their invaluable assistance. TN and MA are JSPS fellows.

REFERENCES

- Ajiki, M., et al. 2003, *AJ*, 126, 2091
- Bertin, E., & Arnouts, S. 1996, *A&AS*, 117, 393
- Bouwens, R. J., et al. 2003, *ApJ*, 595, 589
- Bouwens, R. J., et al. 2004, *ApJ*, in press (astro-ph/0403167)
- Brocklehurst, M. 1971, *MNRAS*, 153, 471
- Bruzual, G., & Charlot, S. 2003, *MNRAS*, 344, 1000
- Bunker, A. J., Stanway, E., Ellis, R. S., & McMahon, R. G. 2004, *MNRAS*, in press (astro-ph/0403223)
- Dickinson, M., et al. 2004, *ApJ*, 600, L99
- Giavalisco, M., et al. 2004, *ApJ*, 600, L103
- Hu, E. M., et al. 2002, *ApJ*, 568, L75; Erratum, *ApJ*, 576, L99
- Hu, E. M., Cowie, L. L., Capak, P., McMahon, R. G., Hayashino, T., & Komiyama, Y. 2004, *AJ*, 127, 563
- Iye, M., et al. 2004, *PASJ*, 56, 381
- Iwata, I., et al. 2003, *PASJ*, 55, 415
- Kaifu, N., et al. 2000, *PASJ*, 52, 1
- Kashikawa, N., et al. 2002, *PASJ*, 54, 819
- Kennicutt, R. C., Jr. 1998, *ARA&A*, 36, 189
- Kodaira, K., et al. 2003, *PASJ*, 55, L17
- Loeb, A., & Barkana, R. 2001, *ARA&A*, 39, 19
- Malhotra, S., & Rhoads, J. E. 2002, *ApJ*, 565, L71
- Miyazaki, S., et al. 2002, *PASJ*, 54, 833
- Ouchi, M., et al. 2003, *ApJ*, 582, 60
- Ouchi, M., et al. 2004, *ApJ*, in press (astro-ph/0309655)
- Schaerer, D. 2000, in *Building the Galaxies: From the Primordial Universe to the Present*, eds. F. Hammer, et al. (Gif-sur-Yvette: Editions Frontières), 389
- Shapley, A. E., Steidel, C. C., Pettini, M., & Adelberger, K. L. 2003, *ApJ*, 588, 65
- Stanway, E., Bunker, A. J., & McMahon, R. G. 2003, *MNRAS*, 342, 439

- Steidel, C. C., Adelberger, K. L. , Giavalisco, M., Dickinson, M., & Pettini, M. 1999, ApJ, 519, 1
- Steidel, C. C., Giavalisco, M., Dickinson, M., & Adelberger, K. L. 1996a, AJ, 112, 352
- Steidel, C. C., Giavalisco, M., Pettini, M., Dickinson, M., & Adelberger, K. L. 1996b, ApJ, 462, L17
- Taniguchi, Y., Shioya, Y., Fujita, S. S., Nagao, T., Murayama, T., & Ajiki, M. 2003, JKAS, 36, 123 (astro-ph/0306409); Erratum, JKAS, 36, 283
- Yagi M., Kashikawa, N., Sekiguchi, M., Doi, M., Yasuda, N., Shimasaku, K. and Okamura, S., 2002, AJ, 123, 66

Table 1. A summary of optical imaging observations

Band	Exposure Time (s)	$m_{\text{AB}}(\text{lim})^{\text{a}}$
B	35700	28.45
V	20400	27.74
R_{c}	36000	27.80
i'	48060	27.43
z'	30240	26.62
$NB816$	36000	26.63
$NB921$	53940	26.54

^aLimiting magnitude (AB) for a 3σ detection on a $2''.0$ diameter aperture.




Fig. 1.— Thumbnail images of SDF J132440.6+273607. The square regions around SDF J132440.6+273607 in the B , V , R_c , i' , $NB816$, z' , and $NB921$ images are shown from left to right. Panel and circle sizes are $16''$ and $8''$, respectively.




Fig. 2.— Sky-subtracted optical position-velocity spectrogram (upper panel) and one-dimensional spectrum (middle panel) of SDF J132440.6+273607 obtained with FOCAS on Subaru. Spectrum of the sky emission is also shown in the lower panel. The spatial size of the displayed area in the upper panel is $8''.1$.




Fig. 3.— (*Left*) Color of $z' - \text{NB921}$ of our 48 i' -dropout high- z galaxy candidates as a function of the z' magnitude. The error-bars denote 1σ uncertainties. The data of SDF J132440.6+273607 is marked by a red circle. The NB921-depressed objects with a $> 2\sigma$ significance are labeled from a to g, sorted in order of the z' magnitude. The magenta, green, light blue, and deep blue lines denote 0σ , 1σ , 2σ , and 3σ uncertainties of the color of $z' - \text{NB921}$, respectively. The lower direction in this diagram means a depression of the NB921 flux compared to the z' flux. (*Right*) Same as the left panel but shown as a function of $i' - z'$ color. For the data with only upper limited $i' - z'$ colors, an arrow is given. The NB921-depressed objects with a $> 2\sigma$ significance are labeled also in this panel. The green, blue and magenta lines denote the expected color of model galaxies with $EW(\text{Ly}\alpha)_0 = 65, 130, \text{ and } 260 \text{ \AA}$, respectively.